

Registration No.:

--	--	--	--	--	--	--	--	--	--

Total Number of Pages: 02

Course: B.Tech  
Sub\_Code: RMA3A001

3<sup>rd</sup> Semester Back Examination: 2025-26

SUBJECT: MATHEMATICS - III

BRANCH(S): AE, AEIE, AERO, AME, AUTO, BIOTECH, CIVIL, CSE, CSEAI, CSEAIML, CSEDS, CSIT, CST, ECE, EEE, ELECTRICAL, ELECTRICAL & C.E, ELECTRONICS & C.E, ETC, IT, MANUTECH, MECH, METTA, MINERAL, MINING, MME, MMEAM, PLASTIC

Time: 3 Hours

Max Marks: 100

Q.Code: U496

Answer Q1 (Part-I) which is compulsory, any eight from Part-II and any two from Part-III.

The figures in the right-hand margin indicate marks.

Part-I

Q1 Answer the following questions: (2 x 10)

- Write the iterative formula for the Newton-Raphson method to find a root of  $f(x) = 0$ .
- Discuss the difference between Doolittle method and Crout method?
- What is the geometrical interpretation of the Trapezoidal Rule?
- Using modified Euler method, find  $y(0.6)$  if  $h = 0.3$  and  $y' = x^2 + y^2, y(0) = 1$
- Define conditional probability with a suitable example.
- State the Baye's theorem.
- If  $X$  follows a Binomial Distribution  $B(n, p)$ , write the expressions for its Mean and Variance.
- Write 2 properties of correlation coefficients.
- Define the probability density function of a Normal Distribution with mean  $\mu$  and variance  $\sigma^2$ .
- Differentiate between a Null Hypothesis ( $H_0$ ) and an Alternative Hypothesis ( $H_1$ ).

Part-II

Q2 Only Focused-Short Answer Type Questions- (Answer Any Eight out of Twelve) (6 x 8)

- How many iterations is required to approximation a root  $f(x) = 0$  in  $[a, b]$ , for which error less than  $10^{-5}$  in bisection method?
- Using Newton's divided difference interpolation, find  $y(10)$  given that  $y(5) = 12, y(6) = 13, y(9) = 14, y(11) = 16$ .
- Solve the system of equations using Crout's Method (LU Decomposition):  $x + y + z = 1, 3x + y - 3z = 5, x - 2y - 5z = 10$ .
- Evaluate  $\int_0^6 \frac{1}{1+x^2} dx$  using Simpson's  $\frac{3}{8}$  Rule.
- Using the Runge-Kutta method of order two, find  $y(0.6)$  if  $h = 0.1$  for  $y' = x^2 + y^2, y(0) = 1$ .
- Evaluate  $\int_0^1 (1 + \frac{\sin x}{x}) dx$  correct to 3 decimal places using trapezoidal rule.

g) A bag contains 4 white and 6 black balls. Two balls are drawn at random without replacement. Find the probability that both balls are black.

h) A random variable  $X$  has the following probability distribution:

$X$	0	1	2	3	4
$P(X)$	$k$	$3k$	$5k$	$7k$	$9k$

Then find the value of  $k$  and calculate  $P(X \geq 2)$ .

i) If 3 % of the electric bulbs manufactured by a company are defective, find the probability that in a sample of 100 bulbs, exactly 4 bulbs are defectives. (Use Poisson Distribution)

j) In a normal distribution, 31 % of the items are under 45 and 8 % are over 64. Find the mean and standard deviation of the distribution.

k) Calculate the Coefficient of Correlation between  $X$  and  $Y$  from the following data:

$X$	1	2	3	4	5
$Y$	2	5	11	8	14

l) Find the Maximum Likelihood Estimator (MLE) for the parameters Binomial distribution.

### Part-III

#### Only Long Answer Type Questions (Answer Any Two out of Four)

**Q3** a) Solve the system of linear equations  $10x + 2y + z = 9, 2x + 20y - 2z = -44, -2x + 3y + 10z = 22$  using the Gauss-Seidel Method. Perform 3 iterations starting with initial guess  $(0,0,0)$ . **(8 x 2)**

b) Using Newton-Raphson method find the smallest positive real root of the equation  $x^4 - 3x^2 + x - 10 = 0$ .

**Q4** a) Apply Euler's Modified Method to find the value of  $y(0.2)$  given  $\frac{dy}{dx} = x + \sqrt{y}$ ,  $y(0) = 1$ , taking step size  $h = 0.1$ . **(8 x 2)**

b) Evaluate the integral  $\int_0^1 e^{-x^2}$  by using Gauss-Legendre 2-point integration formula.

**Q5** a) The joint probability density function of two random variables  $X$  and  $Y$  is given by  $f(x,y) = c(xy + e^x)$  for  $0 < x < 1, 0 < y < 1$  and 0 otherwise. Find the value of  $c$  and the marginal density functions of  $X$  and  $Y$ . **(8 x 2)**

b) State and prove Chebyshev's Inequality for a random variable  $X$  with mean  $\mu$  and variance  $\sigma^2$ .

**Q6** a) Compute and interpret the correlation coefficient for the following data of 6 students selected at random: **(8 x 2)**

Mathematics grade	70	92	80	74	65	83
English grade	74	84	63	87	78	90

b) Obtain the lines of regression for the following data and estimate  $Y$  when  $X = 10$

$X$	1	2	3	4	5
$Y$	3	4	5	7	11